

INFORMED CONSENT FOR ROOT CANAL TREATMENT

Patient's Name	Date
PLEASE INITIAL EACH PARAGRAPH AFTER R	READING. IF YOU HAVE ANY
QUESTIONS, PLEASE ASK YOUR DOCTOR DU	
Only the RCT is to be performed at this office. The performent, etc.) will be done by my regular dentist.	manent outside restoration (filling, onlay,
I acknowledge full responsibility for the payment of suc AT or BEFORE COMPLETION of treatment.	ch services and agree to pay for them in full
Success rate - Although Root Canal therapy has a very high procedure, so it cannot be guaranteed. Root canal treatment (RCT) after proper final restoration by your general dentist (e.g., crown, in Canal therapy may require retreatment, surgery, or even extraction.	can fail but is usually about 95% successful alay, etc.). Occasionally, a tooth which has Root
I have the choice of EXTRACTION, doing nothing (underst treatment at all), or retreatment if RCT had been done previously.	anding infection can occur, if there is no
The tooth/teeth may still need surgery or extraction after the	root canal treatment.
Loss of feeling (i.e., numb ness) or tingling in the lip, tongue temporarily occur, (rarely may be permanent)	e, chin, gums, cheeks, and teeth, may
Canals can be calcified (blocked) and may require root end sinfection) is present or develops.	surgery, if pain persists or root pathology (e.g.,
Complications resulting from use of dental instruments may side of the root, while attempting to find canals) and instruments seend surgery necessary which can affect the prognosis.	
Swelling, sensitivity, pain and infection are possible during a	and after RCT (antibiotics may be needed.)
I may be given injections of local anesthetics. Occasionally needle can injure a nerve to my tongue or lip leading to temporary, the path of the nerve.	
Reactions to anesthetic injections may result in swelling, allo discoloration of skin of face).	ergic reaction or hematoma (temporary
The tooth may be fractured, but not detectable (causing personal content of the c	istent biting pain)
Jaw cramps and muscle spasms can occur (i.e., TMJ difficul-	ty).

Damage to bridges, existing fillings, crowns, chipping of obtaining access to the canals may occur. The tooth and/or roots of tooth. If no crown is present, one usually should be made with	(s) may crack after RCT, possibly resulting in loss
Referred pain to ear, neck and head is possible for 2 to 3	days after RCT.
Antibiotics may cause diarrhea, abdominal cramps, colitication of any of the these complications.	is and allergic reactions (hives). CALL to notify the
It is not unusual to have biting sensitivity up to 4 weeks	after RCT.
Temporary drowsiness/ lack of coordination from oral semay occur.	edative medications/nitrous oxide (laughing gas)
Antibiotics may reduce effectiveness of birth control pill	s (inform doctor).
Minor burn of lip, tongue, or cheek from heated instrume	ent may occur.
No guarantee or assurance can be given to me that the proto my complete satisfaction. Due to individual patient difference re-treatment, or worsening of my present condition despite the contract that therapy would be helpful, and that a worsening of my present recommended treatment (possibly leading to infection).	es, there exists a risk of failure, relapse, selective care provided. However, it is the doctor's opinion
I certify that I speak, read and write English and have rea	ad and fully understand this consent for treatment.
PLEASE ASK YOUR DOCTOR, IF Y	YOU HAVE QUESTIONS
CONCERNING THIS CON	NSENT FORM.
Patient's (or Legal Guardian's) Signature	Date
Doctor's Signature	Date
Witness' Signature	Date